

A PROOF OF QUARK CONFINEMENT IN QCD *

Kei-Ichi Kondo

Department of Physics, Faculty of Science, Chiba University, Chiba 263, JAPAN

ABSTRACT

I propose to reformulate the gauge field theory as the perturbative deformation of a novel topological quantum field theory. It is shown that this reformulation leads to quark confinement in QCD₄. Similarly, the fractional charge confinement is also derived in the strong coupling phase of QED₄. As a confinement criterion, we use the area decay of the expectation value of the Wilson loop.

1. Introduction

An idea of magnetic monopole and its condensation was used in the analytical proofs of "quark" confinement so far:

1975: Polyakov¹ for 3D compact U(1) gauge theory

1977: Polyakov² for 3D Georgi-Glashow model

1994: Seiberg-Witten³ for 4D N=2 SUSY YM/QCD

How to prove quark confinement in QCD₄ analytically? Recent numerical simulations have established that also in QCD the magnetic monopole plays the dominant role in quark confinement (monopole dominance) under the Abelian projection.⁴ To explain this fact analytically and then to prove quark confinement, we propose to use a novel type of topological quantum field theory (TQFT)⁵ which describes 4D magnetic monopole, and to reformulate QCD as a perturbative deformation of the TQFT.

We can show:⁶⁻⁹

1. The TQFT is derived from QCD (without any approximation) in the maximal Abelian gauge (MAG).
2. The calculation of the Wilson loop in 4D YM theory is reduced to that in 2D nonlinear sigma model (NLSM) due to the dimensional reduction. This is an exact result.
3. Area decay of the Wilson loop is derived from 2D Instanton calculus in NLSM.

*Report-no: CHIBA-EP-107 (hep-th/9808186); Talk given at the 3rd International conference on Quark Confinement and the Hadron Spectrum, 7-12 June 1998, Jefferson Lab., Newport News, VA, USA.

2. Basic Idea

We take into account the topological nontrivial background field Ω_μ which corresponds to 4D magnetic monopole current k_μ . The current k_μ obeys the topological conservation law $\partial_\mu k_\mu = 0$ and hence k_μ denotes a loop in 4D spacetime. Usually, Ω_μ denotes an arbitrary, but fixed background (as a solution of classical field equation). In our case, we sum up all topological nontrivial background Ω_μ after integrating out quantum fluctuations Q_μ ,

$$Z_{QCD} = \sum_{\Omega} \int [dQ_\mu] \exp\{-S_{QCD}[\Omega_\mu + Q_\mu]\}, \quad (1)$$

where

$$S_{QCD} = \int d^D x \left[-\frac{1}{2g^2} \text{tr}_G(\mathcal{F}_{\mu\nu} \mathcal{F}_{\mu\nu}) + \bar{\psi}(i\gamma^\mu \mathcal{D}_\mu[\mathcal{A}] - m)\psi \right]. \quad (2)$$

We must regard this background to be different from the 4D instanton discussed in the topological YM theory (Witten⁵) and also from 3D magnetic monopole (Polyakov²).

3. Quantization of YM theory based on BRST formalism

Using the nilpotent BRST transformation δ_B , it is well known that the gauge fixing and the Faddeev-Popov (FP) ghost part is written in the form,

$$\mathcal{L}_{gf} = -i\delta_B G_{gf}, \quad G_{gf} := \text{tr}_G \left\{ \bar{C} \left(F[\mathcal{A}] + \frac{\alpha}{2} \phi \right) \right\}, \quad (3)$$

where α is the gauge-fixing parameter and ϕ is the Lagrange-multiplier field. For the manifestly covariant Lorentz gauge, we choose $F[\mathcal{A}] = \partial^\mu \mathcal{A}_\mu$.

In the maximal Abelian gauge (MAG),^{6,7} a suitable choice of gauge fixing parameter $\alpha = -2$ leads to

$$G_{gf} := \bar{\delta}_B \text{tr}_{G/H} \left(\frac{1}{2} \mathcal{A}_\mu(x) \mathcal{A}_\mu(x) + iC(x) \bar{C}(x) \right). \quad (4)$$

where $\bar{\delta}_B$ is the (nilpotent) anti-BRST transformation. Using the decomposition,

$$\mathcal{A}_\mu(x) = \Omega_\mu(x) + Q_\mu(x) = \frac{i}{g} U(x) \partial_\mu U(x)^\dagger + U(x) \mathcal{V}_\mu(x) U^\dagger(x), \quad (5)$$

the action of the TQFT is given by

$$S_{TQFT} := \int d^D x \, i\delta_B \bar{\delta}_B \text{tr}_{G/H} \left(\frac{1}{2} \Omega_\mu(x) \Omega_\mu(x) + iC(x) \bar{C}(x) \right), \quad (6)$$

and the partition function is given by

$$Z_{TQFT} = \int [dU][dC][d\bar{C}][d\phi] \exp\{-S_{TQFT}[\Omega_\mu]\}, \quad (7)$$

where U is an element of the gauge group G .

4. Strategy of a proof

We consider D-dim. QCD (with a gauge group G) for $D > 2$.

4.1. Step 1:⁷

In MAG, D-dim. QCD (QCD_D) is reformulated as a perturbative deformation of TQFT_D . MAG is a partial gauge fixing which fixes the coset part G/H and leaves the maximal torus group H invariant.

4.2. Step 2:⁷

It is shown that TQFT_D is equivalent to the coset G/H nonlinear sigma model (NLSM) in $(D-2)$ dimensions. This is a consequence of Parisi-Sourlas dimensional reduction due to the hidden supersymmetry in TQFT (6). As extensively discussed more than 20 years ago, QCD_4 and NLSM_2 have common properties: renormalizability, asymptotic freedom ($\beta(g) < 0$), dynamical mass generation, existence of instanton solution, no phase transition for any value of coupling constant (one phase), etc. This similarity between two theories can be understood from this correspondence,

$$\text{QCD}_4 \supset \text{TQFT}_4 \iff G/H \text{ NLSM}_2 \quad (8)$$

For $G = SU(2)$, G/H NLSM is equivalent to $O(3)$ NLSM. Existence of 2D instanton is guaranteed for any N , because $\Pi_2(SU(N)/U(1)^{N-1}) = \Pi_1(U(1)^{N-1}) = \mathbf{Z}^{N-1}$.

4.3. Step 3:⁹

We define the full string tension σ by

$$\sigma := - \lim_{A(C) \rightarrow \infty} \frac{1}{A(C)} \ln \langle W^C[\mathcal{A}] \rangle_{YM_4}, \quad (9)$$

where $W^C[\mathcal{A}]$ is the non-Abelian Wilson loop with the area $A(C)$,

$$W^C[\mathcal{A}] := \text{tr} \left[\mathcal{P} \exp \left(iq \oint_C \mathcal{A}_\mu^A(x) T^A dx^\mu \right) \right]. \quad (10)$$

On the other hand, the diagonal string tension σ_{Abel} is defined by

$$\sigma_{Abel} := - \lim_{A(C) \rightarrow \infty} \frac{1}{A(C)} \ln \langle W^C[a^\Omega] \rangle_{TFT_4}, \quad (11)$$

using the diagonal Wilson loop,

$$W^C[a^\Omega] = \exp \left(iq \oint_C dx^\mu a_\mu^\Omega(x) \right), \quad a_\mu^\Omega(x) := \Omega_\mu^3(x) := \text{tr}(T^3 \Omega_\mu(x)). \quad (12)$$

From the dimensional reduction, we find

$$\langle W^C[a^\Omega] \rangle_{TFT_4} = \langle W^C[a^\Omega] \rangle_{\text{NLSM}_2}. \quad (13)$$

Then it is shown that in the large Wilson loop limit

$$|\sigma - \sigma_{Abel}| \downarrow 0 \quad (A(C) \uparrow \infty). \quad (14)$$

This is derived from the non-Abelian Stokes theorem. For the large (non-intersecting planar) Wilson loop, the full string tension σ is saturated by the diagonal string tension σ_{Abel} . This explains the Abelian dominance and magnetic monopole dominance.


4.4. Step 4:⁷

The whole problem is reduced to calculating

$$\langle W^C[a^\Omega] \rangle_{NLSM_2} = \langle e^{i\frac{q}{g}8\pi Q_S} \rangle_{NLSM_2}, \quad Q_S = \frac{1}{8\pi} \int_S d^2z \, \epsilon_{\mu\nu} \mathbf{n} \cdot (\partial_\mu \mathbf{n} \times \partial_\nu \mathbf{n}). \quad (15)$$

Note that the integrand of Q_S is the instanton density in $NLSM_2$. Therefore, Q_S counts the number of instantons minus that of anti-instantons inside the area S bounded by the Wilson loop C . In order to perform the actual calculation, we adopt the naive instanton calculus (dilute gas approximation). This leads to the area law of the diagonal Wilson loop and the non-zero diagonal string tension σ_{Abel} for the fractional charge q (i.e., when q/g is not an integer). More systematic instanton calculations enable us to identify the instanton solution with the Coulomb gas of vortices.

Note that all the above steps are exact except for the instanton calculus of the Wilson loop in $NLSM$. The above results imply that the non-zero string tension σ in QCD_4 follows from the non-zero diagonal string tension σ_A in $NLSM_2$. The problem of proving area law in QCD_4 is reduced to the equivalent problem in $NLSM_2$.



Similar strategy is also applied to QED_4 to prove the existence of strong coupling confinement phase.⁸ This follows from the existence of Berezinski-Kosterlitz-Thouless transition of the $O(2)$ $NLSM_2$.

References

1. A.M. Polyakov, Phys. Lett. B **59**, 82 (1975).
2. A.M. Polyakov, Nucl. Phys. B **120**, 429 (1977).
3. N. Seiberg and E. Witten, Nucl. Phys. B **426**, 19 (1994); **431**, 484 (1994).
4. G. 't Hooft, Nucl. Phys. B **190** [FS3], 455 (1981).
5. E. Witten, Commun. Math. Phys. **117**, 353 (1988).
6. K.-I. Kondo, hep-th/9709109 (revised), Phys. Rev. D **57**, 7467 (1998).
K.-I. Kondo, hep-th/9803063, Prog. Theor. Phys. Suppl., No. 130, in press.
7. K.-I. Kondo, hep-th/9801024 (revised), Phys. Rev. D **58** (1998), in press.
8. K.-I. Kondo, hep-th/9803133 (revised), Phys. Rev. D **58** (1998), in press.
9. K.-I. Kondo, hep-th/9805153 (revised), Phys. Rev. D **58** (1998), in press.